



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁷ : C12N 15/10</p>	<p>A1</p>	<p>(11) International Publication Number: WO 00/09680 (43) International Publication Date: 24 February 2000 (24.02.00)</p>
<p>(21) International Application Number: PCT/US99/17405 (22) International Filing Date: 30 July 1999 (30.07.99) (30) Priority Data: 60/096,009 10 August 1998 (10.08.98) US 09/363,209 29 July 1999 (29.07.99) US (71) Applicant: CHIRON CORPORATION [US/US]; 4560 Horton Street, Emeryville, CA 94608 (US). (72) Inventors: MITTELSTAEDT, Denise, M.; 13687 Quinton Road, San Diego, CA 92129 (US). HSU, David, Chi-Tang; 7387 Mannix Court, San Diego, CA 92109 (US). (74) Agents: MCMASTERS, David, D. et al.; Seed and Berry LLP, 6300 Columbia, 701 Fifth Avenue, Seattle, WA 98104-7092 (US).</p>		<p>(81) Designated States: CA, JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published With international search report.</p>
<p>(54) Title: METHODS FOR PURIFYING NON-CHROMOSOMAL NUCLEIC ACID MOLECULES FROM CELLS</p> <div data-bbox="324 1144 1299 1428"> </div> <p>(57) Abstract</p> <p>Methods are provided for purifying non-chromosomal nucleic acid molecules from cells, comprising: the general steps of (a) lysing cells to form a nucleic acid-containing lysate, and (b) applying the lysate to a depth filter in order to obtain a clarified solution containing purified non-chromosomal nucleic acid molecules.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

METHODS FOR PURIFYING NON-CHROMOSOMAL NUCLEIC ACID MOLECULES FROM CELLS

TECHNICAL FIELD

The present invention relates generally to biotechnology, and more specifically, to a method for purifying non-chromosomal nucleic acid molecules from a cell lysate.

BACKGROUND OF THE INVENTION

Since the discovery of the structure of nucleic acid molecules (DNA and RNA) in the 1950's, research into the structure, function, and use of these molecules has increased dramatically. In particular, a tremendous amount of research has been undertaken in order to utilize these molecules to diagnose and treat disease. In order however to obtain sufficient quantities of nucleic acids for use in experiments, and for diagnosis and treatment of disease, it is first necessary to purify and or isolate substantial quantities of nucleic acid molecules. For example, in the field of gene therapy, patients can be vaccinated or treated with nucleic acid molecules in order to protect against, or, remedy a disease (*see, e.g.*, PCT publication WO 95/07994). However, preparation of large-quantities of purified, pharmaceutical-grade nucleic acid molecules is presently time-consuming and costly.

More specifically, while many methods exist for the purification of nucleic acid molecules, such methods are often limited when the nucleic acid molecules are to be produced for therapeutic purposes, since they must be prepared free of any contaminants such as toxic compounds and antigenic molecules. Prior art methods of DNA purification often utilized highly toxic chemicals such as ethidium bromide, phenol, chloroform, and cesium chloride (Sambrook, et al., 1992, *Molecular Cloning: A Laboratory Manual* (2nd ed.), Cold Spring Harbor, NY). Additionally, many methods involving both physical or chemical lysis often relied upon centrifugation to remove cellular debris, a step which is undesirable because of the difficulty in maintaining sterility. Furthermore, procedures using alkali lysis (Birboim and Doly, *Nucleic Acids*

Res. 7;1513, 1979) may lead to the loss of a significant amount of DNA through degradation caused by the increase in pH.

The present invention provides methods for producing purified nucleic acid molecules that are suitable for use in pharmaceutical applications, and on a large scale. Such methods do not require traditional techniques such as toxic extractants, mutagenic reagents, or steps such as centrifugation. The present invention also provides other, related advantages.

SUMMARY OF THE INVENTION

Briefly stated, the present invention provides methods for purifying non-chromosomal nucleic acid molecules from cells, comprising the general steps of (a) lysing cells to form a nucleic acid-molecule containing lysate, (b) precipitating chromosomal DNA and cell wall components from the lysate, in order to form a flocculent solution, and (c) applying the flocculent solution to a depth filter, such that non-chromosomal nucleic acid molecules are purified.

Within further embodiments, the non-chromosomal nucleic acid molecules can be further isolated by chromatographically separating non-chromosomal nucleic acid molecules (*e.g.*, by ion-exchange chromatography or preparative HPLC) from proteins and other compounds which are in the lysate. Within other embodiments, the clarified solution may be concentrated (*e.g.*, by centrifugation or tangential flow). Within yet other embodiments the purification and/or isolation methods described herein can be performed in a sterile environment and/or in a continuous manner.

The methods provided herein may be utilized to purify and/or isolate a variety of non-chromosomal nucleic acid molecules, including for example plasmids and RNA.

Within certain embodiments of the invention, the depth filter has a nominal pore size of between about 0.2 and 20 microns. Within other embodiments, the non-chromosomal nucleic acid molecules are precipitated utilizing a neutralizing agent such as sodium acetate or potassium acetate.

The methods provided herein may be utilized for purifying and/or isolating non-chromosomal nucleic acid molecules from a wide variety of cells,

including for example, bacterial cells, yeast cells, and mammalian cells (*i.e.*, both eukaryotic and non-karyotic cells).

These and other aspects of the present invention will become evident upon reference to the following detailed description and attached drawings. In addition, various references are set forth herein which describe in more detail certain procedures or compositions (*e.g.*, plasmids, etc.), and are therefore incorporated by reference in their entirety.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a schematic illustration of one representative purification method of the present invention.

Figure 2 is a photograph which shows the removal of precipitated debris.

DETAILED DESCRIPTION OF THE INVENTION

DEFINITIONS

Prior to setting forth the invention, it may be helpful to an understanding thereof to first set forth definitions of certain terms that will be used hereinafter.

"Depth filter" refers to a filter which is capable excluding particles of a selected size from a given media. Depth filters, (as opposed to membrane filters) exclude particles from media by capturing particles within the matrix of the media (as compared to primarily the surface of the media for membrane filters). Depth filters are typically composed of fibrous media, usually cellulose, glass fiber, or, polypropylene (or a combination of these).

"Purified" non-chromosomal nucleic acid molecules, as utilized within the context of the present invention, refers to nucleic acid molecules that are substantially free of cellular debris and chromosomal DNA (*i.e.*, has an absorbance of 0.010 as determined by spectrophotometric analysis at 600 nm, and preferably less than 0.005, 0.002, or, 0.001).

"Isolated" non-chromosomal nucleic acid molecules, as utilized within the context of the present invention, refers to a preparation of nucleic acid molecules

which, when linearized and applied to an agarose gel, upon application of an appropriate voltage form a single broad band of nucleic acid molecules. If the nucleic acid molecules are not linearized, a broad band or 3 separate close bands of supercoiled (dimers or oligomers), relaxed and/or linear nucleic acids are obtained. Protein assays can also be performed in order to ensure that the protein content is less than 10 ug/ml, and preferably less than 5, 2, or 1 ug/ml.

As noted above, the present invention provides methods for purifying non-chromosomal nucleic acid molecules from cells, comprising the general steps of (a) lysing cells to form a nucleic acid-molecule containing lysate, (b) precipitating chromosomal DNA and cell wall components from the lysate, in order to form a flocculent solution, and (c) applying the flocculent solution to a depth filter, such that non-chromosomal nucleic acid molecules are purified. Briefly, by utilizing a depth filter for removing cellular debris and other precipitates that are formed due to the lysing of cells, one can readily purify non-chromosomal nucleic acid molecules without the need for traditional separation steps such as centrifugation, or filtration through membrane filters such as Watman paper, or, cheesecloth. Such methods may be readily accomplished without the use of polyethylene glycol, or organic extractants (*e.g.*, phenol or ethidium bromide).

In order to further the understanding of the invention, set forth below is a brief discussion of methods for lysing cells, precipitating chromosomal DNA and cell wall components, and filtering the resultant flocculent solution. Furthermore, methods are provided for further isolating and/or concentrating the purified non-chromosomal nucleic acid molecules. Also, a discussion is provided on the use of these non-chromosomal molecules in research and pharmaceutical applications.

LYSING CELLS

Utilizing the methods of the present invention, non-chromosomal nucleic acid molecules can be purified from a wide variety of cells, including for example both mammalian (*e.g.*, human, rat and mouse) and non-mammalian (*e.g.*, bacterial, yeast, or insect) cells. Briefly, the first step is to lyse the cells in order to form a lysate containing cytoplasmic proteins and plasmids. A wide variety of methods

can be utilized to lyse cells, including for example, physical methods such as homogenization, sonication or the use of a French Press, and chemical (*e.g.*, chemical lysis) and enzymatic methods (*e.g.*, lysozyme).

Briefly, in order to homogenize cells, cellular material is first blended with a mortar and pestle or a blender. Typically for purification of non-chromosomal nucleic acid molecules from bacteria, a bacterial pellet is resuspended into a solution and a mortar and pestle or blender is utilized to rupture cell walls. One alternative method of homogenization is the French Press. Briefly, within such a press a cell pellet is suspended in solution and squeezed between two hard surfaces. Once the cell wall is ruptured, cellular components containing cytoplasmic proteins and plasmids are released.

In order to sonicate cells, cells are placed in a sonication bath, or alternatively, a probe-type oscillator is placed into a vessel containing the cells. Representative examples of suitable sonicators include the Sonifier II® Cell Disrupter (Branson), or, the Ultrasonic Cleaner (Branson).

Chemical lysis may readily be accomplished by a variety of methods. For example, within one embodiment a bacterial pellet is suspended in buffer. The solution is rapidly and completely mixed with 0.2M NaOH and 1% SDS (or equivalent). This is allowed to incubate for a maximum of five minutes. The solution is then neutralized with either potassium or sodium acetate. This forms a precipitate consisting of cellular debris and genomic DNA. It is also possible during this step to introduce RNase A for reduction of free RNA.

Cells may also be lysed by enzymatic means utilizing commonly available enzymes such as lysozyme.

Any of the above procedures can be utilized to lyse cells in order to form a lysate containing cytoplasmic proteins and plasmids. Briefly, these processes result in the protein or plasmid being released from the cell primarily causing large holes in the cell wall. The cellular debris and chromosomal DNA trapped within the cell form a precipitate. Separation of this precipitate can occur by different methods dependent on the material required.

PRECIPITATING A LYSATE FROM A FLOCCULENT SOLUTION

A wide variety of methods may be utilized to precipitate a lysate (e.g., cellular debris and chromosomal DNA) from a flocculent solution, including for example, chemically (e.g., by pH change or the addition of salt), or physically (e.g., by the use of heat). As an illustration, during chemical lysis, high pH of the lysate can be achieved by the addition of sodium hydroxide to the lysis buffer, followed by sodium acetate or potassium acetate to neutralize the lysate, and to cause precipitation of cellular debris and chromosomal DNA. If homogenization is utilized to lyse cells, either streptomycin sulfate or ammonium sulfate may be utilized to precipitate the lysate.

Utilizing the methods provided herein, at least 5 to 6 liters of flocculent solution may be processed utilizing just one filter with ten square feet of surface area. In order to scale up the processing of fluids, multiple filters may be added in order to process in parallel (or, alternatively, serially) larger volumes of fluids.

FILTERING A LYSATE

As noted above, once a flocculent solution has been generated, the solution is applied to a depth filter. As noted above, the term 'depth filter' refers to a filter which are capable excluding particles of a selected size from a given media. Depth filters, (as opposed to membrane filters) exclude particles from media by capturing particles within the matrix of the media (as compared to primarily the surface of the media for membrane filters).

Depth filters are typically composed of fibrous media, usually cellulose, glass fiber, or, polypropylene (or a combination of these). Depth filters suitable for use within the present invention may be purchased commercially, at, for example, Sartorius Corporation (Edgewood, N.Y.), Millipore, or Gillman. Depth filters are also described within U.S. Patent No. 4,594,202 (Method of Making Cylindrical Fibrous filter Structures"), U.S. Patent No. 4,726,901 (entitled "Cylindrical Fibrous Structures With Graded Pore Size"), and EPO 0148638 entitled "Cylindrical Fibrous Structures and Method of Manufacture").

Utilizing such techniques, non-chromosomal nucleic acid molecules can be readily purified from at least 10, 15, or even 25 grams (wet weight) per liter of cells, at a substantially faster rate than conventional methods. Moreover, the process described herein is scalable, allowing 1, 2, 3, 4, 5 liters (or even greater quantities) of cells to be lysed and filtered for non-chromosomal nucleic acid molecules. For example, 125 grams of cells (wet weight) in 5 liters can be filtered using a depth filter with 5-10 square feet of surface area.

ISOLATING AND/OR CONCENTRATING NON-CHROMOSOMAL NUCLEIC ACID MOLECULES

After the flocculent solution is applied to a depth filter, the resultant solution containing purified non-chromosomal nucleic acid molecules may, optionally, be applied to further process steps in order to isolate and or further concentrate the non-chromosomal nucleic acid molecules.

For example, utilizing chromatographic techniques, (e.g., ion exchange and/or size exclusion) one can readily isolate the non-chromosomal nucleic acid molecules. Briefly, ion exchange utilizes anion columns including either DEAE or Q resins. Qiagen, Santa Clarita has a patent for use of DEAE resin in plasmid purification while the primary Q resin patent for use in plasmid DNA is by Biosepra, Boston. Experiments with alternative resins have shown consistency in the results. There is also an affinity resin available from Puresyn, Boston that can be used after ion exchange. Typically the concentration of the plasmid DNA following chromatography is dilute.

Concentration of the purified and/or isolated non-chromosomal nucleic acids may also be accomplished utilizing techniques such as precipitation and Tangential Flow Filtration.

Briefly, precipitation usually involves the addition of an alcohol (e.g., isopropanol or ethanol) to the solution followed by centrifugation. The plasmid DNA forms a pellet that is then rinsed with ethanol and allowed to air dry. The plasmid DNA can then be dissolved into any buffer.

Tangential flow filtration can be used by limiting the cut-off size and both reducing the volume, removing small particle contaminants and acting as a buffer

exchange. With Tangential Flow Filtration, the cut-off size required is much smaller than expected due to the compact properties of supercoiled plasmid.

METHODS FOR UTILIZING PURIFIED OR ISOLATED NUCLEIC ACIDS, AND
PHARMACEUTICAL COMPOSITIONS

Purified or isolated nucleic acid molecules of the present invention have a variety of uses. For example, within one embodiment of the invention the isolated nucleic acid molecules may be utilized for diagnostic assays, or, alternatively, for therapeutic purposes.

When utilized for therapeutic purpose, the purified and/or isolated nucleic acid molecules may be further formulated with one or more excipients, buffers or stabilizers. In addition, the nucleic acid molecules will typically be provided in a sterile, pyrogen-free form, in a suitable dosage for administration for the intended purposes.

Representative examples of such therapeutic techniques and methods are provided in PCT Publication Nos. WO 95/07994 and WO 96/40952 and US. Patent Nos. 5,580,859 and 5,589,466.

The following examples are offered by way of illustration, and not by way of limitation.

EXAMPLES

EXAMPLE 1

PURIFICATION OF PLASMID DNA FROM *E. COLI* DH5- α CELLS

This example demonstrates a method of purifying plasmids from *E. coli* DH5- α cells. Briefly, from *E. coli* DH5- α cells containing the ELVS 1.5 β -gal plasmid (see WO 97/38087) were grown in Digene superbrot with phosphate (Digene, Beltsville, Md) containing the antibiotic kanamycin (50 mg/ml, Sigma, St. Louis, Mo). The cells were incubated for 20-28 hours at 37° C in a shaker bath. The cells were then harvested by centrifugation at 5000 x G) for 15 minutes at 2-10° C. The supernatant

was discarded and the pellets were resuspended in 2 liters of P1-RNase-A buffer (50 mM Tris/HCl, 10 mM EDTA, pH 8.0, containing 100 mg RNase-A per liter, Qiagen, Santa Clarita, CA). The cells were then lysed upon addition of 2.15 liters of lysis buffer P2 (200 mM NaOH, 1% SDS, Qiagen, Santa Clarita, CA). This was accomplished by allowing the resuspended cells to come in contact with the lysis buffer through a "Y" connector tube which connected 3 pieces of tubing (see Figure 1). This method of mixing does not require inversion. A dual head peristaltic pump was used to generate the flow of the fluids.

P3 buffer containing 3.0 M potassium acetate pH 5.5 (Qiagen, Santa Clarita, CA) was added to the above lysed cells through a similar type of "Y" connector tube, except that a second pump was used to simultaneously regulate the flow at an approximate 2:1 ratio of lysate to P3 buffer. This solution formed a precipitate and was allowed to incubate on ice for a minimum of 30 minutes.

The precipitated solution was pumped directly through a Sartopure PP2 0.5m depth filter capsule (Sartorius, Inc., Yauca, P.R.). Endotoxin was removed by adding Endotoxin Removal (ER) Buffer (Qiagen, Santa Clarita, CA) to the filtered pool and mixing uniformly.

Following incubation for a minimum of 1 hour on ice, the combined ER buffer and filtered lysate pool was transferred to the chromatography unit, an ion exchange column (Ultrapure 100, Qiagen, Santa Clarita, CA). The run cycle was set at a flow rate of 5 mL/min. for 18 hours prior to washing with QC buffer (1 M NaCl, 50 mM MOPS, 15% Ethanol, pH 7.0) at a flow rate of 20 ml/min for 2 1/2 hours. The plasmids were eluted with QN buffer (1.6 M NaCl, 50 mM MOPS, 15% Ethanol, pH 7.0). The plasmids were then precipitated with isopropanol to a final concentration of 42%, centrifuged for 60 minutes at 13,000 x G at 2-8°C, washed with 70% ethanol, and resuspended in TE (10mM TRIS-HCl, 1mM EDTA). Upon subsequent, analysis utilizing a Pierce Protein Microassay, less than 1.25 ug/ml protein was detected.

From the foregoing, it will be appreciated that, although specific embodiments of the invention have been described herein for purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention. Accordingly, the invention is not limited except as by the appended claims.

CLAIMS

We claim:

1. A method for purifying non-chromosomal nucleic acid molecules from cells, comprising:

- a) lysing cells to form a nucleic acid molecule-containing lysate;
- b) precipitating chromosomal DNA and cell wall components from the lysate, in order to form a flocculent solution; and
- c) applying the flocculent solution to a depth filter, such that non-chromosomal nucleic acid molecules are purified.

2. The method according to claim 1 wherein the step of lysing is accomplished by homogenizing cells.

3. The method according to claim 1 wherein the step of lysing is accomplished by chemical lysis.

4. The method according to claim 1, further comprising the step of chromatographically isolating said purified non-chromosomal nucleic acid molecules.

5. The method according to claim 4 wherein said purified non-chromosomal nucleic acid molecules are separated on an ion-exchange column.

6. The method according to claim 1, further comprising the step of concentrating said purified non-chromosomal nucleic acid molecules.

7. The method according to claim 4 wherein said purified non-chromosomal nucleic acid molecules are concentrated by tangential flow.

8. The method according to claim 1 wherein said non-chromosomal nucleic acid molecules are plasmids.

9. The method according to claim 1 wherein said non-chromosomal nucleic acid molecules are RNA.

10. The method according to claim 1 wherein said depth filter has a nominal pore size of between 0.2 and 20 microns.


11. The method according to claim 1 wherein any one of steps a) through c) are performed in a sterile environment.

12. The method according to claim 1 wherein any one of steps a) through c) are performed in a continuous manner.

13. The method according to claim 1 wherein said cells are bacteria.

14. The method according to claim 1 wherein said cells are eukaryotic cells.

1/2

*Fig. 1*

2/2




Fig. 2

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/17405

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C12N15/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97 29190 A (THEREXSYS LTD) 14 August 1997 (1997-08-14) page 27	1-14
X	WO 93 11218 A (DIAGEN INST MOLEKULARBIO) 10 June 1993 (1993-06-10) page 2-4; claims 1-15; examples 1,2	1-14
A	WO 96 02658 A (LEE ANN L ;MERCK & CO INC (US); SAGAR SANGEETHA (US)) 1 February 1996 (1996-02-01) page 8, line 27 -page 9, line 9; examples 1,5; table 1	1-14
A	US 5 645 986 A (VAZIRI HOMAYOUN ET AL) 8 July 1997 (1997-07-08) page 42, line 50 - line 62	2
	--- -/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

20 October 1999

Date of mailing of the international search report

26/10/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

van Klompenburg, W

INTERNATIONAL SEARCH REPORT

Inter :nal Application No

PCT/US 99/17405

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>US 5 576 196 A (MARQUET MAGDA ET AL) 19 November 1996 (1996-11-19) column 11, line 12 - line 30</p>	7

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/17405

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9729190 A	14-08-1997	AU 1799497 A	28-08-1997
		CA 2244115 A	14-08-1997
		EP 0880585 A	02-12-1998
WO 9311218 A	10-06-1993	DE 4139664 A	03-06-1993
		DE 59205979 D	15-05-1996
		DE 59209549 D	10-12-1998
		WO 9311221 A	10-06-1993
		EP 0616638 A	28-09-1994
		EP 0616639 A	28-09-1994
		EP 0875271 A	04-11-1998
		JP 7501222 T	09-02-1995
		JP 7501223 T	09-02-1995
WO 9602658 A	01-02-1996	AU 708798 B	12-08-1999
		AU 3126295 A	16-02-1996
		CA 2192342 A	01-02-1996
		EP 0771355 A	07-05-1997
		JP 10503086 T	24-03-1998
US 5645986 A	08-07-1997	US 5695932 A	09-12-1997
		US 5489508 A	06-02-1996
		AU 682082 B	18-09-1997
		AU 1209095 A	29-05-1995
		AU 1330795 A	29-05-1995
		AU 6058298 A	04-06-1998
		CA 2173872 A	18-05-1995
		EP 0728207 A	28-08-1996
		JP 2875394 B	31-03-1999
		JP 9502102 T	04-03-1997
		WO 9513381 A	18-05-1995
		WO 9513383 A	18-05-1995
		US 5891639 A	06-04-1999
		US 5629154 A	13-05-1997
		US 5648215 A	15-07-1997
		US 5686306 A	11-11-1997
		US 5639613 A	17-06-1997
		US 5837453 A	17-11-1998
		US 5693474 A	02-12-1997
		US 5863726 A	26-01-1999
		US 5804380 A	08-09-1998
		US 5830644 A	03-11-1998
		AU 688262 B	12-03-1998
		AU 4374093 A	13-12-1993
		AU 7183698 A	20-08-1998
		AU 8949598 A	07-01-1999
		AU 8949698 A	14-01-1999
		CA 2135648 A	25-11-1993
		CA 2245461 A	25-11-1993
		CA 2245462 A	25-11-1993
		EP 0642591 A	15-03-1995
		JP 11123100 A	11-05-1999
		JP 11127874 A	18-05-1999
		JP 8501079 T	06-02-1997
		WO 9323572 A	25-11-1993
		US 5744300 A	28-04-1998
		US 5686245 A	11-11-1997
		US 5840495 A	24-11-1998

INTERNATIONAL SEARCH REPORT

Information on patent family members

Inter: International Application No

PCT/US 99/17405

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5645986 A		US 5707795 A	13-01-1996
US 5576196 A	19-11-1996	CA 2206265 A	18-07-1996
		EP 0802975 A	29-10-1997
		JP 11502403 T	02-03-1999
		WO 9621729 A	18-07-1996